




Restoration and Monitoring of Aquatic Quality in a Coal-Mined Watershed, Swatara Creek at Ravine, Pennsylvania

Charles A. Cravotta III, Michael D. Bilger, Robin A. Brightbill (U.S. Geological Survey, New Cumberland, PA 17070) and Daniel Bogar (Pennsylvania Department of Environmental Protection, Harrisburg, PA 17105)




Swatara Creek drains an area of 577 mi² within the Ridge and Valley Province in east-central Pennsylvania. The upper 43-mi² area above Ravine is underlain by the Southern Anthracite Field. Although several surface and underground anthracite mines presently are active, most mines in the Swatara Creek Basin were abandoned before 1960. Once abandoned, the underground mines flooded producing numerous large discharges that are contaminated with acidity, sulfate, and metals. The polluted drainage from these abandoned mines affects water quality miles downstream. Construction of the proposed Swatara State Park Reservoir has been delayed until the AMD is remediated.







To neutralize AMD in the Swatara Creek Basin above Ravine, a variety of treatment systems was installed in 1995 through 1997 including limestone-sand dosing, open limestone channels, anoxic and oxic limestone drains, limestone diversion wells, and wetlands. Additional treatments and reclamation work have been completed since then. To characterize untreated mine drainage and treatment-system performance, data on flow rate and water quality throughout the basin were collected during baseflow and stormflow conditions in 1995-2001. Data for the station on Swatara Creek, near Ravine, indicate cumulative effects of AMD remediation and transport of pollutants from the mined part of the basin.



Baseflow data collected periodically since 1959 at Ravine indicate significant improvement in water quality. For example, sulfate declined while pH increased sharply after 1975 from 3.5-4.4 (median ~4) to 4.6-7.0 (median ~6). The decline in SO₄ concentration probably was caused by a decline in pyrite oxidation after flooding of the abandoned mines had minimized inflows of oxygenated air and water. The associated increase in pH was caused by the onset of carbonate buffering that occurred when the rate of alkalinity production equaled or exceeded acid production. Although a variety of environmental factors could affect pH and SO₄ concentrations, consistently near-neutral pH with variable SO₄ concentration at Ravine during 1998-2000, particularly during stormflow, imply that the recently implemented limestone treatments have neutralized further improving water quality. At near-neutral pH, the transport of dissolved metals typically is attenuated owing to precipitation and adsorption. Nevertheless, recent monitoring indicates elevated concentrations and transport of Fe, Al, Mn, and trace metals during stormflow and elevated concentrations of Fe, Mn, Co, Cu, Pb, Ni, and Zn in streambed sediments relative to unmined areas and to toxicity guidelines for aquatic invertebrates and fish. The metals are ubiquitous in the fine fraction (<0.063 mm) of bed sediment in mining-affected tributaries and the main stem of Swatara Creek and represent a long-term source of contamination.



As a consequence of the improved water quality in Swatara Creek at Ravine, the fish community has rebounded. No fish were present during ecological surveys in 1985 and 1990; however, in 1994 and 1996 six species of fish were found. Increasing numbers of fish species have been found annually since 1996. In 2000, twenty-four species of fish were documented, including cold-water species such as brook trout and sculpin and warm-water species such as sunfish, pickerel, and bullhead catfish. Although the majority of the fish species are considered to have moderate tolerance to pollution, several intolerant species including river chub, cutlips minnow, and longnose dace, have been reported since 1997.



An increased abundance of benthic macroinvertebrate taxa that are considered intolerant of pollution indicates water quality improved from fair in 1994 to very good in 2000. Nevertheless, Hydropsychidae (caddisflies) and Chironomidae (midges), which are known to tolerate acidic conditions, were dominant. Although subordinate, the appearance of Ephemeroptera (mayflies) in 1997 and later years is significant in that these animals are sensitive to acidic conditions and considered intolerant to pollution. The benthic macroinvertebrate community recorded for 1999 and 2000 can be characterized as moderately impacted based on total taxa and slightly impacted based on total mayfly, stonefly, and caddisfly taxa.

REFERENCES:

Cravotta, C.A., III, and Bilger, M.D., 2001, Water-quality trends for a coal-mined watershed in Eastern Pennsylvania: Geochemistry--Exploration, Environment, Analysis, vol. 1, p. 33-50.

Cravotta, C.A., III, and Weitzel, J.B., 2001, Detecting change in water quality from implementation of limestone treatment systems in a coal-mined watershed, Pennsylvania, in 8th National Nonpoint Source Monitoring Workshop: U.S. Environmental Protection Agency EPA/905-R-01-008.